

Symbolic Software for the Painlevé Test of Nonlinear Ordinary and Partial Differential Equations

Douglas BALDWIN and Willy HEREMAN

Department of Mathematical and Computer Sciences, Colorado School of Mines
Golden, CO 80401, USA

E-mail: painlevetest@douglasbaldwin.com and whereman@mines.edu

Received April 22, 2005; Accepted in Revised Form June 5, 2005

Abstract

The automation of the traditional Painlevé test in *Mathematica* is discussed. The package `PainleveTest.m` allows for the testing of polynomial systems of nonlinear ordinary and partial differential equations which may be parameterized by arbitrary functions (or constants). Except where limited by memory, there is no restriction on the number of independent or dependent variables. The package is quite robust in determining all the possible dominant behaviors of the Laurent series solutions of the differential equation. The omission of valid dominant behaviors is a common problem in many implementations of the Painlevé test, and these omissions often lead to erroneous results. Finally, our package is compared with the other available implementations of the Painlevé test.

1 Introduction

Completely integrable nonlinear partial differential equations (PDEs) have remarkable properties, such as infinitely many generalized symmetries, infinitely many conservation laws, the Painlevé property, Bäcklund and Darboux transformations, bilinear forms, and Lax pairs (cf. [2, 11, 24, 25]). Completely integrable equations model physically interesting wave phenomena in reaction-diffusion systems, population and molecular dynamics, nonlinear networks, chemical reactions, and waves in material science. By investigating the complete integrability of a nonlinear PDE, one gains important insight into the structure of the equation and the nature of its solutions.

Broadly speaking, Painlevé analysis is the study of the singularity structure of differential equations. Specifically, a differential equation is said to have the Painlevé property if all the movable singularities of all its solutions are poles. There is strong evidence [48, 50, 51] that integrability is closely related to the singularity structure of the solutions of a differential equation (cf. [33, 38]). For instance, dense branching of solutions around movable singularities has been shown to indicate nonintegrability [49].

At the turn of the nineteenth-century, Painlevé [30] and his colleagues classified all the rational second-order ODEs for which all the solutions are single-valued around all movable

singularities. Equations possessing this property could either be solved in terms of known functions or transformed into one of the six Painlevé equations whose solutions define the Painlevé transcendents. The Painlevé transcendents cannot be expressed in terms of the classical transcendental functions, except for special values of their parameters [19].

The complex singularity structure of solutions was first used by Kovalevskaya in 1889 to identify a new integrable system of equations for the motion for a rotating top (cf. [14, 38]). Ninety years later, Ablowitz, Ramani and Segur (ARS) [2, 3] and McLeod and Olver [27] formulated the Painlevé conjecture which gives a useful *necessary* condition for determining whether a PDE is solvable using the Inverse Scattering Transform (IST) method. Specifically, the Painlevé conjecture asserts that every nonlinear ODE obtained by an exact reduction of a nonlinear PDE solvable by the IST-method has the Painlevé property. While necessary, the condition is not sufficient; in general, most PDEs do not have exact reductions to nonlinear ODEs and therefore satisfy the conjecture by default [41]. Weiss, Tabor and Carnevale (WTC) [44] proposed an algorithm for testing PDEs directly (which is analogous to the ARS algorithm for testing ODEs). For a thorough discussion of the traditional Painlevé property, see [1, 8, 10, 13, 18, 28, 31, 33, 38, 39].

There are numerous methods for solving completely integrable nonlinear PDEs, for instance by explicit transformations into linear equations or by using the IST-method [11]. Recently, progress has been made using *Mathematica* and *Maple* in applying the IST-method to difficult equations, including the Camassa-Holm equation [21]. While there is as yet no systematic way to determine if a differential equation is solvable using the IST-method [27], having the Painlevé property is a strong indicator that it will be.

There are several implementations of the Painlevé test in various computer algebra systems, including *Reduce*, *MacSyma*, *Maple* and *Mathematica*. The implementations described in [34, 35, 37] are limited to ODEs, while the implementations discussed in [16, 45–47] allow the testing of PDEs directly using the WTC algorithm. The implementation for PDEs written in *Mathematica* by Hereman et al. [16] is limited to two independent variables (x and t) and is unable to find all the dominant behaviors in systems with undetermined exponents α_i (as is the case with the Hirota-Satsuma system). Our package `PainleveTest.m` [4] written in *Mathematica* syntax, allows the testing of polynomial PDEs (and ODEs) with no limitation on the number of differential equations or the number of independent variables (except where limited by memory). Our implementation also allows the testing of differential equations that have undetermined dominant exponents α_i and that are parameterized by arbitrary functions (or constants). The implementations for PDEs written in *Maple* by Xu and Li [45–47] were written after the one presented in this paper and are comparable to our implementation.

The paper is organized as follows: in Section 2 we review the basics of Painlevé analysis. Section 3 discusses the WTC algorithm for testing PDEs and uses the Korteweg-de Vries (KdV) equation and the Hirota-Satsuma system of coupled KdV (cKdV) equations to show the subtleties of the algorithm. We detail the algorithms to determine the dominant behavior, resonances, and constants of integration using a generalized system of coupled nonlinear Schrödinger (NLS) equations in Section 4. Additional examples are presented in Section 5 to illustrate the capabilities of the software. Section 6 compares our software package to other codes and briefly discusses the generalizations of the WTC algorithm. The use of the package `PainleveTest.m` [4] is shown in Section 7. We draw some conclusions and discuss the results in Section 8.

2 Painlevé Analysis

Consider a system of M polynomial differential equations,

$$F_i(\mathbf{u}(\mathbf{z}), \mathbf{u}'(\mathbf{z}), \mathbf{u}''(\mathbf{z}), \dots, \mathbf{u}^{(m_i)}(\mathbf{z})) = \mathbf{0}, \quad i = 1, 2, \dots, M, \quad (2.1)$$

where the dependent variable $\mathbf{u}(\mathbf{z})$ has components $u_1(\mathbf{z}), \dots, u_M(\mathbf{z})$, the independent variable \mathbf{z} has components z_1, \dots, z_N , and $\mathbf{u}^{(k)}(\mathbf{z}) = \partial^k \mathbf{u}(\mathbf{z}) / (\partial z_1^{k_1} \partial z_2^{k_2} \dots \partial z_N^{k_N})$ denotes the collection of mixed derivative terms of order k . Let $m = \sum_{i=1}^M m_i$, where m_i is the highest order in each equation. If there are any arbitrary coefficients (constants or analytic functions of \mathbf{z}) parameterizing the system, we assume they are nonzero. For simplicity, in the examples we denote the components of $\mathbf{u}(\mathbf{z})$ by $u(\mathbf{z}), v(\mathbf{z}), w(\mathbf{z}), \dots$, and the components of \mathbf{z} by x, y, z, \dots, t .

A differential equation has the *Painlevé property* if all the movable singularities of all its solutions are poles. A singularity is *movable* if it depends on the constants of integration of the differential equation. For instance, the Riccati equation,

$$w'(z) + w^2(z) = 0, \quad (2.2)$$

has the general solution $w(z) = 1/(z - c)$, where c is the constant of integration. Hence, (2.2) has a movable simple pole at $z = c$ because it depends on the constant of integration. Solutions of ODEs can have various kinds of singularities, including branch points and essential singularities; examples of the various types of singularities [23] are shown in Table 1. As a general property, solutions of *linear* ODEs have only fixed singularities [19].

Simple <i>fixed</i> pole	$z w' + w = 0$	\Rightarrow	$w(z) = c/z$
Simple <i>movable</i> pole	$w' + w^2 = 0$	\Rightarrow	$w(z) = 1/(z - c)$
Movable <i>algebraic branch</i> point	$2ww' - 1 = 0$	\Rightarrow	$w(z) = \sqrt{z - c}$
Movable <i>logarithmic branch</i> point	$w'' + w'^2 = 0$	\Rightarrow	$w(z) = \log(z - c_1) + c_2$
Non-isolated movable <i>essential</i> singularity	$(1 + w^2)w'' + (1 - 2w)w'^2 = 0$	\Rightarrow	$w(z) = \tan\{\ln(c_1 z + c_2)\}$

Table 1. Examples of various types of singularities.

In general, a function of several complex variables cannot have an isolated singularity [29]. For example, $f(z) = 1/z$ has an isolated singularity at the point $z = 0$, but the function $f(w, z) = 1/z$ of two complex variables, $w = u + iv, z = x + iy$, has a two-dimensional manifold of singularities, namely the points $(u, v, 0, 0)$, in the four-dimensional space of these variables. Therefore, we will define a pole of a function of several complex variables as a point (a_1, a_2, \dots, a_N) , in whose neighborhood the function can be written

in the form $f(\mathbf{z}) = h(\mathbf{z})/g(\mathbf{z})$, where g and h are both analytic in a region containing (a_1, \dots, a_N) in its interior, $g(a_1, \dots, a_N) = 0$, and $h(a_1, \dots, a_N) \neq 0$.

The WTC algorithm considers the singularity structure of the solutions around non-characteristic manifolds of the form $g(\mathbf{z}) = 0$, where $g(\mathbf{z})$ is an analytic function of $\mathbf{z} = (z_1, z_2, \dots, z_N)$ in a neighborhood of the manifold. Specifically, if the singularity manifold is determined by $g(\mathbf{z}) = 0$ and $\mathbf{u}(\mathbf{z})$ is a solution of the PDE, then one assumes a Laurent series solution

$$u_i(\mathbf{z}) = g^{\alpha_i}(\mathbf{z}) \sum_{k=0}^{\infty} u_{i,k}(\mathbf{z}) g^k(\mathbf{z}), \quad i = 1, 2, \dots, M, \quad (2.3)$$

where the coefficients $u_{i,k}(\mathbf{z})$ are analytic functions of \mathbf{z} with $u_{i,0}(\mathbf{z}) \neq 0$ in a neighborhood of the manifold and the α_i are integers with at least one exponent $\alpha_i < 0$. The requirement that the manifold $g(\mathbf{z}) = 0$ is non-characteristic, ensures that the expansion (2.3) is well defined in the sense of the Cauchy-Kovalevskaya theorem [41, 43].

Substituting (2.3) into (2.1) and equating coefficients of like powers of $g(\mathbf{z})$ determines the possible values of α_i and defines a recursion relation for $u_{i,k}(\mathbf{z})$. The recursion relation is of the form

$$Q_k \mathbf{u}_k = \mathbf{G}_k(\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{k-1}, g, \mathbf{z}), \quad \mathbf{u}_k = (u_{1,k}, u_{2,k}, \dots, u_{M,k})^T, \quad (2.4)$$

where Q_k is an $M \times M$ matrix and T denotes transpose.

For (2.1) to pass the Painlevé test, the series (2.3) should have $m - 1$ arbitrary functions as required by the Cauchy-Kovalevskaya theorem (as $g(\mathbf{z})$ is the m -th arbitrary function). If so, the Laurent series solution corresponds to the general solution of the equation [1]. The $m - 1$ arbitrary functions $u_{i,k}(\mathbf{z})$ occur when k is one of the roots of $\det(Q_k)$. These roots $r_1 \leq r_2 \leq \dots \leq r_m$ are called *resonances*. The resonances are also equal to the Fuchs indices of the auxiliary equations of Darboux [7].

Since the WTC algorithm is unable to detect essential singularities, it is only a necessary condition for the PDE to have the Painlevé property [6]. While rarely done in practice, sufficiency is proved by finding a transformation which linearizes the differential equation, yields an auto-Bäcklund transformation, a Bäcklund transformation, or hodographic transformation [15] to another differential equation which has the Painlevé property (see [8, 23, 33] for more information).

3 Algorithm for the Painlevé Test

In this section, we outline the WTC algorithm for testing PDEs for the Painlevé property. We discuss the Kruskal simplification and the Painlevé test of ODEs after the three main steps are outlined. Each of these steps is illustrated using both the KdV equation and the cKdV equations due to Hirota and Satsuma. Details of the three main steps of the algorithm are postponed till Section 4.

Step 1 (Determine the dominant behavior). It is sufficient to substitute

$$u_i(\mathbf{z}) = \chi_i g^{\alpha_i}(\mathbf{z}), \quad i = 1, 2, \dots, M, \quad (3.1)$$

where χ_i is a constant, into (2.1) to determine the leading exponents α_i . In the resulting polynomial system, equating every two or more possible lowest exponents of $g(\mathbf{z})$ in each

equation gives a linear system for α_i . The linear system is then solved for α_i and each solution branch is investigated. The traditional Painlevé test requires that all the α_i are integers and that at least one is negative.

If any of the α_i are non-integer in a given branch, then that branch of the algorithm terminates. A non-integer α_i implies that some solutions of (2.1) have movable algebraic branch points. Often, a suitable change of variables in (2.1) can remove the algebraic branch point. An alternative approach is to use the “weak” Painlevé test, which allows certain rational α_i and resonances; see [13, 18, 32, 33] for more information.

If one or more α_i remain undetermined, we assign integer values to the free α_i so that every equation in (2.1) has at least two different terms with equal lowest exponents.

For each solution α_i , we substitute

$$u_i(\mathbf{z}) = u_{i,0}(\mathbf{z})g^{\alpha_i}(\mathbf{z}), \quad i = 1, 2, \dots, M, \quad (3.2)$$

into (2.1). We then solve the (typically) nonlinear equation for $u_{i,0}(\mathbf{z})$, which is found by balancing the leading terms. By leading terms, we mean those terms with the lowest exponent of $g(\mathbf{z})$. If any of the solutions contradict the assumption that $u_{i,0}(\mathbf{z}) \neq 0$, then that branch of the algorithm fails the Painlevé test.

If any of the α_i are non-integer, all the α_i are positive, or there is a contradiction with the assumption that $u_{i,0}(\mathbf{z}) \neq 0$, then that branch of the algorithm terminates and does not pass the Painlevé test for that branch.

Step 2 (Determine the resonances). For each α_i and $u_{i,0}(\mathbf{z})$, we calculate the $r_1 \leq \dots \leq r_m$ for which $u_{i,r}(\mathbf{z})$ is an arbitrary function in (2.3). To do this, we substitute

$$u_i(\mathbf{z}) = u_{i,0}(\mathbf{z})g^{\alpha_i}(\mathbf{z}) + u_{i,r}(\mathbf{z})g^{\alpha_i+r}(\mathbf{z}) \quad (3.3)$$

into (2.1), and keep only the lowest order terms in $g(\mathbf{z})$ that are linear in $u_{i,r}$. This is done by computing the solutions for r of $\det(Q_r) = 0$, where the $M \times M$ matrix Q_r satisfies

$$Q_r \mathbf{u}_r = \mathbf{0}, \quad \mathbf{u}_r = (u_{1,r} \ u_{2,r} \ \dots \ u_{M,r})^T. \quad (3.4)$$

If any of the resonances are non-integer, then the Laurent series solutions of (2.1) have a movable algebraic branch point and the algorithm terminates. If r_m is not a positive integer, then the algorithm terminates; if $r_1 = -1, r_2 = \dots = r_m = 0$ and $m - 1$ of the $u_{i,0}(\mathbf{z})$ found in Step 1 are arbitrary, then (2.1) passes the Painlevé test. If (2.1) is parameterized, the values for $r_1 \leq \dots \leq r_m$ may depend on the parameters, and hence restrict the allowable values for the parameters.

There is always a resonance $r = -1$ which corresponds to the arbitrariness of $g(\mathbf{z})$; as such, it is often called the universal resonance. When there are negative resonances other than $r = -1$, (or, more than one resonance equals -1) then the Laurent series solution is not the general solution and further analysis is needed to determine if (2.1) passes the Painlevé test. The perturbative Painlevé approach, developed by Conte et al. [9], is one method for investigating negative resonances.

Step 3 (Find the constants of integration and check compatibility conditions). For the system to possess the Painlevé property, the arbitrariness of $u_{i,r}(\mathbf{z})$ must be verified up to the highest resonance level. This is done by substituting

$$u_i(\mathbf{z}) = g^{\alpha_i}(\mathbf{z}) \sum_{k=0}^{r_m} u_{i,k}(\mathbf{z})g^k(\mathbf{z}) \quad (3.5)$$

into (2.1), where r_m is the largest positive integer resonance.

For (2.1) to have the Painlevé property, the $(M + 1) \times M$ augmented matrix $(Q_k | \mathbf{G}_k)$ must have rank M when $k \neq r$ and rank $M - s$ when $k = r$, where s is the algebraic multiplicity of r in $\det(Q_r) = 0$, $1 \leq k \leq r_m$, and Q_k and \mathbf{G}_k are as defined in (2.4). If the augmented matrix $(Q_k | \mathbf{G}_k)$ has the correct rank, solve the linear system (2.4) for $u_{1,k}(\mathbf{z}), \dots, u_{M,k}(\mathbf{z})$ and use the results in the linear system at level $k + 1$.

If the linear system (2.4) does not have a solution, then the Laurent series solution of (2.1) has a movable logarithmic branch point and the algorithm terminates. Often, when (2.1) is parameterized, carefully choosing the parameters will resolve the difference in the ranks of Q_k and $(Q_k | \mathbf{G}_k)$.

If the algorithm does not terminate, then the Laurent series solutions of (2.1) are free of movable algebraic or logarithmic branch points and (2.1) passes the Painlevé test.

The Painlevé test of PDEs is quite cumbersome; in particular, Step 3 is lengthy and prone to error when done by hand. To simplify Step 3, Kruskal proposed a simplification which now bears his name. In the context of the WTC algorithm, it is sometimes called the Weiss-Kruskal simplification [20, 23]. The manifold defined by $g(\mathbf{z}) = 0$ is non-characteristic, that means $g_{z_l}(\mathbf{z}) \neq 0$ for some l on the manifold $g(\mathbf{z}) = 0$. By the implicit function theorem, we can then locally solve $g(\mathbf{z}) = 0$ for z_l , so that

$$g(\mathbf{z}) = z_l - h(z_1, \dots, z_{l-1}, z_{l+1}, \dots, z_N), \quad (3.6)$$

for some arbitrary function h . Using (3.6) greatly simplifies the computation of the constants of integration $u_{i,k}(z_1, \dots, z_{l-1}, z_{l+1}, \dots, z_N)$. However, with the Kruskal simplification one loses the ability to use the Weiss truncation method [42] to find a linearising transformation, an auto-Bäcklund transformation, or a Bäcklund transformation (see [8]).

When testing ODEs, (2.3) must be replaced by

$$u_i(z) = g^{\alpha_i}(z) \sum_{k=0}^{\infty} u_{i,k} g^k(z), \quad i = 1, 2, \dots, M, \quad (3.7)$$

where the coefficients $u_{i,k}$ are constants, $g(z) = z - z_0$, and z_0 is an arbitrary constant. If z explicitly occurs in the ODE, then it is (automatically) replaced by $g(z) + z_0$ prior to Step 1 of the test. An example of the Painlevé test of an ODE is given in Section 5.

3.1 The Korteweg-de Vries equation

To illustrate the steps of the algorithm, let us examine the KdV equation [1],

$$u_t + 6uu_x + u_{3x} = 0, \quad (3.8)$$

the most famous completely integrable PDE from soliton theory. Note that for simplicity, we use $u_{ix} = u_{xx \dots x} = \partial^i u / \partial x^i$ and $g_{ix} = \partial^i g / \partial x^i$ when $i \geq 3$.

Substituting (3.1) into (3.8) gives

$$\alpha \chi \{ 6t g^{\alpha-1} + 6\chi g_x g^{2\alpha-1} + g^{\alpha-3} [(\alpha-1)((\alpha-2)g_x^2 + 3gg_{xx})g_x + g^2 g_{3x}] \} = 0. \quad (3.9)$$

The lowest exponents of $g(x, t)$ are $\alpha - 3$ and $2\alpha - 1$. Equating these leading exponents gives $\alpha = -2$. Substituting (3.2), $u(x, t) = u_0(x, t)g^{-2}(x, t)$, into (3.8) and requiring that the leading terms (in $g^{-5}(x, t)$) balance, gives $u_0(x, t) = -2g_x^2(x, t)$.

Substituting (3.3), $u(x, t) = -2g_x^2(x, t)g^{-2}(x, t) + u_r(x, t)g^{r-2}(x, t)$, into (3.8) and equating the coefficients of the dominant terms (in $g^{r-5}(x, t)$) that are linear in $u_r(x, t)$ gives

$$(r - 6)(r - 4)(r + 1)g_x(x, t)^3 = 0. \quad (3.10)$$

Assuming $g_x(x, t) \neq 0$, the resonances of (3.8) are $r_1 = -1, r_2 = 4$ and $r_3 = 6$.

We now substitute

$$\begin{aligned} u(x, t) &= g^{-2}(x, t) \sum_{k=0}^6 u_k(x, t)g^k(x, t) \\ &= -2g_x^2(x, t)g^{-2}(x, t) + u_1(x, t)g^{-1}(x, t) + \cdots + u_6(x, t)g^4(x, t) \end{aligned} \quad (3.11)$$

into (3.8) and group the terms of like powers of $g(x, t)$. So, we will pull off the coefficients of $g^{k-5}(x, t)$ at level k . Equating the coefficients of $g^{-4}(x, t)$ to zero at level $k = 1$, gives $u_1(x, t)g_x^3(x, t) = 2g_x^3(x, t)g_{xx}(x, t)$. Setting $u_1(x, t) = 2g_{xx}(x, t)$, we get

$$u_2(x, t) = -\frac{g_t g_x^2 + 3g_x g_{xx}^2 - 4g_x^2 g_{3x}}{6g_x^3}, \quad (3.12)$$

at level $k = 2$. Similarly, at level $k = 3$,

$$u_3(x, t) = \frac{g_x^2 g_{xt} - g_t g_x g_{xx} + 3g_x^3 - 4g_x g_{xx} g_{3x} + g_x^2 g_{4x}}{6g_x^4}. \quad (3.13)$$

At level $k = r_2 = 4$, we find

$$(u_1)_t + 6\{u_3(u_0)_x + u_2(u_1)_x + u_1(u_2)_x + u_0(u_3)_x\} + (u_1)_{3x} = 0, \quad (3.14)$$

which is trivially satisfied upon substitution of the solutions of $u_0(x, t), \dots, u_3(x, t)$. Therefore, the compatibility condition at level $k = r_2 = 4$ is satisfied and $u_4(x, t)$ is indeed arbitrary. At level $k = 5$, $u_5(x, t)$ is unambiguously determined, but not shown due to length. Finally, the compatibility condition at level $k = r_3 = 6$ is trivially satisfied when the solutions for $u_0(x, t), \dots, u_3(x, t)$ and $u_5(x, t)$ are substituted into the recursion relation at that resonance level.

Therefore, the Laurent series solution $u(x, t)$ of (3.8) in the neighborhood of $g(x, t) = 0$ is free of algebraic and logarithmic movable branch points. Furthermore, since the Laurent series solution,

$$u(x, t) = g^{-2}(x, t) \sum_{k=0}^{\infty} u_k(x, t)g^k(x, t), \quad (3.15)$$

has three arbitrary functions, $g(x, t), u_4(x, t)$, and $u_6(x, t)$, (as required by the Cauchy-Kovalevskaya theorem since (3.8) is of third order) it is also the general solution. Hence, we conclude that (3.8) passes the Painlevé test.

The Weiss-Kruskal simplification uses $g(x, t) = x - h(t)$. Consequently, $g_x = 1, g_{xx} = g_{3x} = \cdots = 0$, and the Laurent series,

$$u(x, t) = g^{-2}(x, t) \sum_{k=0}^{\infty} u_{i,k}(t)g^k(x, t), \quad (3.16)$$

becomes

$$u(x, t) = -\frac{2}{(x - h(t))^2} + \frac{1}{6}h'(t) + u_4(t)(x - h(t))^2 + \frac{1}{36}h''(t)(x - h(t))^3 + u_6(t)(x - h(t))^4 + \dots, \quad (3.17)$$

where $h(t)$, $u_4(t)$ and $u_6(t)$ are arbitrary.

3.2 The Hirota-Satsuma system

To show the subtleties in determining the dominant behavior, consider the cKdV equations due to Hirota and Satsuma [1] with real parameter a ,

$$\begin{aligned} u_t &= a(6uu_x + u_{3x}) - 2vv_x, \quad a > 0, \\ v_t &= -3uv_x - v_{3x}. \end{aligned} \quad (3.18)$$

Again, we substitute (3.1), $u(x, t) = \chi_1 g^{\alpha_1}(x, t)$ and $v(x, t) = \chi_2 g^{\alpha_2}(x, t)$, into (3.18) and pull off the lowest exponents of $g(x, t)$. From the first equation, we get $\alpha_1 - 3$, $2\alpha_1 - 1$, and $2\alpha_2 - 1$. From the second equation, we get $\alpha_2 - 3$ and $\alpha_1 + \alpha_2 - 1$. Hence, $\alpha_1 = -2$ from the second equation. Substituting this into the first equation gives $\alpha_2 \geq -2$.

Substituting (3.2) into (3.18) and requiring that at least two leading terms balance gives us two branches: $\alpha_1 = \alpha_2 = -2$ and $\alpha_1 = -2, \alpha_2 = -1$. The branches with $\alpha_1 = -2$ and $\alpha_2 \geq 0$ are excluded for they require that either $u_0(x, t)$ or $v_0(x, t)$ is identically zero.

Continuing with the two branches and solving for $u_0(x, t)$ and $v_0(x, t)$ gives

$$\begin{cases} \alpha_1 = \alpha_2 = -2, \\ u_0(x, t) = -4g_x^2(x, t), \\ v_0(x, t) = \pm 2\sqrt{6}ag_x^2(x, t), \end{cases} \quad \text{and} \quad \begin{cases} \alpha_1 = -2, \alpha_2 = -1, \\ u_0(x, t) = -2g_x^2(x, t), \\ v_0(x, t) \text{ arbitrary.} \end{cases} \quad (3.19)$$

For the branch with $\alpha_1 = \alpha_2 = -2$, substituting (3.3),

$$\begin{cases} u(x, t) = -4g_x^2(x, t)g^{-2}(x, t) + u_r(x, t)g^{r-2}(x, t), \\ v(x, t) = \pm 2\sqrt{6}ag_x^2(x, t)g^{-2}(x, t) + v_r(x, t)g^{r-2}(x, t), \end{cases} \quad (3.20)$$

into (3.18) and equating to zero the coefficients of the lowest order terms in $g(x, t)$ that are linear in u_r and v_r gives

$$\begin{pmatrix} -(r-4)(r^2-5r-18)ag_x^3(x, t) & \pm 12\sqrt{6}ag_x^3(x, t) \\ \mp 4(r-4)\sqrt{6}ag_x^3(x, t) & (r-2)(r-7)rg_x^3(x, t) \end{pmatrix} \begin{pmatrix} u_r(x, t) \\ v_r(x, t) \end{pmatrix} = \mathbf{0}. \quad (3.21)$$

From

$$\det(Q_r) = -a(r+2)(r+1)(r-3)(r-4)(r-6)(r-8)g_x^6(x, t) = 0, \quad (3.22)$$

we obtain the resonances $r_1 = -2, r_2 = -1, r_3 = 3, r_4 = 4, r_5 = 6$, and $r_6 = 8$.

By convention, the resonance $r_1 = -2$ is ignored since it violates the hypothesis that $g(x, t)^{-2}$ is the dominant term in the expansion near $g(\mathbf{z}) = 0$. Furthermore, this is not a principal branch since the series has only five arbitrary functions instead of the required six

(as the term corresponding to resonance $r_1 = -2$ does not contribute to the expansion). Thus, this leads to a particular solution and the general solution may still be multi-valued.

As in the previous example, the constants of integration at level k are found by substituting (3.5) into (3.18) and pulling off the coefficients of $g^{k-5}(x, t)$. At level $k = 1$,

$$\begin{pmatrix} 11ag_x^3(x, t) & \pm 2\sqrt{6a}g_x^3(x, t) \\ \pm 2\sqrt{6a}g_x^3(x, t) & -g_x^3(x, t) \end{pmatrix} \begin{pmatrix} u_1(x, t) \\ v_1(x, t) \end{pmatrix} = \begin{pmatrix} 20ag_x^3(x, t)g_{xx}(x, t) \\ \pm 10\sqrt{6a}g_x^3(x, t)g_{xx}(x, t) \end{pmatrix}, \quad (3.23)$$

and thus,

$$u_1(x, t) = 4g_{xx}(x, t), \quad v_1(x, t) = \pm 2\sqrt{6a}g_{xx}(x, t). \quad (3.24)$$

At level $k = 2$,

$$\begin{aligned} u_2(x, t) &= \frac{3g_{xx}^2(x, t) - g_x(x, t)(g_t(x, t) + 4g_{3x}(x, t))}{3g_x^2(x, t)}, \\ v_2(x, t) &= \pm \frac{(1 + 2a)g_t(x, t)g_x(x, t) + 4ag_x(x, t)g_{3x}(x, t) - 3ag_{xx}^2(x, t)}{\sqrt{6a}g_x^2(x, t)}. \end{aligned} \quad (3.25)$$

The compatibility conditions at levels $k = r_3 = 3$ and $k = r_4 = 4$ are trivially satisfied. At levels $k = 5$ and $k = 7$, $u_k(x, t)$ and $v_k(x, t)$ are unambiguously determined (not shown). At resonance levels $k = r_5 = 6$ and $k = r_6 = 8$, the compatibility conditions require $a = \frac{1}{2}$.

Likewise, for the branch with $\alpha_1 = -2, \alpha_2 = -1$, substituting (3.3),

$$\begin{cases} u(x, t) = -2g_x^2(x, t)g^{-2}(x, t) + u_r(x, t)g^{r-2}(x, t), \\ v(x, t) = v_0(x, t)g^{-1}(x, t) + v_r(x, t)g^{r-1}(x, t), \end{cases} \quad (3.26)$$

into (3.18) gives

$$\begin{pmatrix} -a(r+1)(r-4)(r-6)g_x^3(x, t) & -3v_0(x, t)g_x(x, t) \\ 0 & r(r-1)(r-5)g_x^3(x, t) \end{pmatrix} \begin{pmatrix} u_r(x, t) \\ v_r(x, t) \end{pmatrix} = \mathbf{0}. \quad (3.27)$$

Since

$$\det(Q_r) = -a(r+1)r(r-1)(r-4)(r-5)(r-6)g_x^6(x, t), \quad (3.28)$$

the resonances are $r_1 = -1, r_2 = 0, r_3 = 1, r_4 = 4, r_5 = 5$, and $r_6 = 6$.

Since $r_2 = 0$ is a resonance, there must be freedom at level $k = r_2 = 0$; indeed, coefficient $u_0(x, t) = -2g_x^2(x, t)$ is unambiguously determined but $v_0(x, t)$ is arbitrary. Then, the constants of integration are found by substituting

$$\begin{cases} u(x, t) = -2g_x^2(x, t)g^{-2}(x, t) + u_1(x, t)g^{-1}(x, t) + \cdots + u_6(x, t)g^4(x, t), \\ v(x, t) = v_0(x, t)g^{-1}(x, t) + v_1(x, t) + \cdots + v_6(x, t)g^5(x, t), \end{cases} \quad (3.29)$$

into (3.18) and pulling off the coefficients of $g^{k-5}(x, t)$ in the first equation and $g^{k-4}(x, t)$ in the second equation. At level $k = r_3 = 1$,

$$\begin{pmatrix} a & 0 \\ v_0(x, t) & 0 \end{pmatrix} \begin{pmatrix} u_1(x, t) \\ v_1(x, t) \end{pmatrix} = \begin{pmatrix} 2ag_{xx}(x, t) \\ 2v_0(x, t)g_{xx}(x, t) \end{pmatrix}. \quad (3.30)$$

So, $u_1(x, t) = 2g_{xx}(x, t)$ and $v_1(x, t)$ is arbitrary. At level $k = 2$,

$$\begin{pmatrix} 12ag_x^2 & 0 \\ -3v_0g_x & -6g_x^3 \end{pmatrix} \begin{pmatrix} u_2 \\ v_2 \end{pmatrix} = \begin{pmatrix} 2g_tg_x + 6ag_{xx}^2 - v_0^2 - 8ag_xg_{3x} \\ v_0g_t + 6(v_1)_xg_x^2 - 3(v_0)_xg_{xx} + 3(v_0)_{2x}g_x + v_0g_{3x} \end{pmatrix}, \quad (3.31)$$

which unambiguously determines $u_2(x, t)$ and $v_2(x, t)$. Similarly, the coefficients in the Laurent series solution are unambiguously determined at level $k = 3$. At resonance level $k = r_4 = 4$, the compatibility condition is trivially satisfied. At resonance levels $k = r_5 = 5$ and $k = r_6 = 6$, the compatibility conditions requires $a = \frac{1}{2}$.

Therefore, (3.18) satisfies the necessary conditions for having the Painlevé property when $a = \frac{1}{2}$, a fact confirmed by other analyses of complete integrability [1].

4 Key Algorithms

In this section, we present the three key algorithms in greater detail. To illustrate the steps we consider a generalization

$$\begin{aligned} iu_t + u_{xx} + (|u|^2 + \beta|v|^2)u + a(x, t)u + c(x, t)v &= 0, \\ iv_t + v_{xx} + (|v|^2 + \beta|u|^2)v + b(x, t)v + d(x, t)u &= 0, \end{aligned} \quad (4.1)$$

of the coupled nonlinear Schrödinger (NLS) equations [40]

$$\begin{aligned} iu_t + u_{xx} + (|u|^2 + \beta|v|^2)u &= 0, \\ iv_t + v_{xx} + (|v|^2 + \beta|u|^2)v &= 0. \end{aligned} \quad (4.2)$$

In (4.1), $a(x, t), \dots, d(x, t)$ are arbitrary complex functions and β is a real constant parameter. Since all the functions in (4.1) are complex, we write the system as

$$\begin{aligned} iu_t + u_{xx} + (u\bar{u} + \beta v\bar{v})u + a(x, t)u + c(x, t)v &= 0, \\ i\bar{u}_t - \bar{u}_{xx} - (u\bar{u} + \beta v\bar{v})\bar{u} - \bar{a}(x, t)\bar{u} - \bar{c}(x, t)\bar{v} &= 0, \\ iv_t + v_{xx} + (v\bar{v} + \beta u\bar{u})v + b(x, t)v + d(x, t)u &= 0, \\ i\bar{v}_t - \bar{v}_{xx} - (v\bar{v} + \beta u\bar{u})\bar{v} - \bar{b}(x, t)\bar{v} - \bar{d}(x, t)\bar{u} &= 0, \end{aligned} \quad (4.3)$$

and treat u, \bar{u}, v , and \bar{v} as independent complex functions. As is customary, the variables with overbars denote complex conjugates.

4.1 Algorithm to determine the dominant behavior

Determining the dominant behavior of (2.1) is delicate and the omission of valid dominant behaviors often leads to erroneous results [33].

Step 1 (Substitute the leading-order ansatz). To determine the values of α_i , it is sufficient to substitute $u_i(\mathbf{z}) = \chi_i g(\mathbf{z})^{\alpha_i}$, into (2.1), where χ_i is constant and $g(\mathbf{z})$ is an analytic function in a neighborhood of the non-characteristic manifold defined by $g(\mathbf{z}) = 0$.

Step 2 (Collect exponents and prune non-dominant branches). The balance of exponents must come from different terms in (2.1). For each equation $F_i = 0$, collect the exponents of $g(\mathbf{z})$. Then, remove non-dominant exponents and duplicates (that come from the same term

in (2.1)). For example, $\alpha_1 + 1$ is non-dominant and can be removed from $\{\alpha_1 - 1, \alpha_1 + 1\}$ since $\alpha_1 - 1 < \alpha_1 + 1$.

For (4.3), the exponents corresponding to each equation are

$$\begin{aligned} F_1 : & \quad \{\alpha_1 - 2, 2\alpha_1 + \alpha_2, \alpha_1 + \alpha_3 + \alpha_4\}, \\ F_2 : & \quad \{\alpha_2 - 2, \alpha_1 + 2\alpha_2, \alpha_2 + \alpha_3 + \alpha_4\}, \\ F_3 : & \quad \{\alpha_3 - 2, 2\alpha_3 + \alpha_4, \alpha_1 + \alpha_2 + \alpha_3\}, \\ F_4 : & \quad \{\alpha_4 - 2, \alpha_3 + 2\alpha_4, \alpha_1 + \alpha_2 + \alpha_4\}, \end{aligned} \tag{4.4}$$

after duplicates and non-dominant exponents have been removed.

Step 3 (Combine expressions and compute relations for α_i). For each F_i separately, equate all possible combinations of two elements. Then, construct relations between the α_i by solving for α_1, α_2 , etc., one at a time.

For (4.4), we get

$$\begin{aligned} F_1 : \quad & \{\alpha_1 - 2 = 2\alpha_1 + \alpha_2, \alpha_1 - 2 = \alpha_1 + \alpha_3 + \alpha_4, \\ & \quad 2\alpha_1 + \alpha_2 = \alpha_1 + \alpha_3 + \alpha_4\} \\ \Rightarrow & \{\alpha_1 + \alpha_2 = -2, \alpha_3 + \alpha_4 = -2, \alpha_1 + \alpha_2 = \alpha_3 + \alpha_4\}. \end{aligned} \tag{4.5}$$

For F_2, F_3 and F_4 we again find that $\{\alpha_1 + \alpha_2 = -2, \alpha_3 + \alpha_4 = -2, \alpha_1 + \alpha_2 = \alpha_3 + \alpha_4\}$.

Step 4 (Combine equations and solve for exponents α_i). By combining the sets of expressions in an ‘‘outer product’’ fashion, we generate all the possible linear equations for α_i . Solving these linear systems, we form a set of all possible solutions for α_i .

For (4.3), we have three sets of linear equations

$$\left\{ \alpha_1 + \alpha_2 = -2 \right. \Rightarrow \left\{ \begin{array}{l} \alpha_1 + \alpha_2 = -2, \\ \alpha_3 + \alpha_4 \geq -2, \end{array} \right. \tag{4.6}$$

$$\left\{ \alpha_3 + \alpha_4 = -2 \right. \Rightarrow \left\{ \begin{array}{l} \alpha_1 + \alpha_2 \geq -2, \\ \alpha_3 + \alpha_4 = -2, \end{array} \right. \tag{4.7}$$

and

$$\left\{ \alpha_1 + \alpha_2 = \alpha_3 + \alpha_4, \right. \Rightarrow \left\{ \alpha_1 + \alpha_2 = \alpha_3 + \alpha_4 \geq -2. \right. \tag{4.8}$$

Although the algorithm treats u, \bar{u}, v , and \bar{v} as independent complex functions, we know that $\alpha_1 = \alpha_2$ and $\alpha_3 = \alpha_4$ because \bar{u} and \bar{v} are the complex conjugates of u and v . Our package `PainleveTest.m` can take advantage of such additional information by using the option `DominantBehaviorConstraints -> {alpha[1] == alpha[2], alpha[3] == alpha[4]}`. Using this additional information yields three cases, $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 \geq -1$ and $\alpha_1 = \alpha_2 = -1, \alpha_3 = \alpha_4 \geq -1$ and $\alpha_1 = \alpha_2 \geq -1, \alpha_3 = \alpha_4 = -1$.

Step 5 (Fix the undetermined α_i). First, compute the minimum values for the undetermined α_i . If a minimum value cannot be determined, then the user-defined value `DominantBehaviorMin` is used. If so, the value of the free α_i is counted up to a user defined `DominantBehaviorMax`. If neither of the bounds is set, the software will run the

test for the default values $\alpha_i = -1, -2$ and -3 . For maximal flexibility, with the option `DominantBehavior` one can also run the code for user-specified values of α_i . An example is given in Section 7. In any case, the selected or given dominant behaviors are checked for consistency with (2.1).

For (4.3), if we take $\alpha_1, \dots, \alpha_4 < 0$, then we are left with only one branch

$$\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = -1. \quad (4.9)$$

Step 6 (Compute the first terms in the Laurent series). Using the values for α_i , substitute

$$u_i(\mathbf{z}) = u_{i,0}(\mathbf{z})g^{\alpha_i}(\mathbf{z}) \quad (4.10)$$

into (2.1) and solve the resulting (typically) nonlinear equations for $u_{i,0}(\mathbf{z})$ using the assumption that $u_{i,0}(\mathbf{z}) \neq 0$.

For (4.3), we find

$$\begin{cases} \alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = -1, \\ u_0(x, t) = -2g_x^2(x, t)(1 + \beta)^{-1}\bar{u}_0^{-1}(x, t), \\ v_0(x, t) = -2g_x^2(x, t)(1 + \beta)^{-1}\bar{v}_0^{-1}(x, t), \end{cases} \quad (4.11)$$

where $\bar{u}_0(x, t)$ and $\bar{v}_0(x, t)$ are arbitrary functions.

If we do not restrict $\alpha_1, \dots, \alpha_4 < 0$, then there are contradictions with the assumption $u_{i,0}(\mathbf{z}) \neq 0$ for all but two possible dominant behaviors,

$$\begin{cases} \alpha_1 = \alpha_2 = -1, \\ \alpha_3 \geq 3, \\ \alpha_4 \geq 3, \end{cases} \quad \text{and} \quad \begin{cases} \alpha_1 \geq 3, \\ \alpha_2 \geq 3, \\ \alpha_3 = \alpha_4 = -1. \end{cases} \quad (4.12)$$

4.2 Algorithm to determine the resonances

Step 1 (Construct matrix Q_r). Substitute

$$u_i(\mathbf{z}) = u_{i,0}(\mathbf{z})g^{\alpha_i}(\mathbf{z}) + u_{i,r}(\mathbf{z})g^{\alpha_i+r}(\mathbf{z}) \quad (4.13)$$

into (2.1). Then, the (i, j) -th entry of the $M \times M$ matrix Q_r is the coefficients of the linear terms in $u_{j,r}(\mathbf{z})$ of the leading terms in equation $F_i = 0$.

Step 2 (Find the roots of $\det(Q_r)$). The resonances are the solutions of $\det(Q_r) = 0$. If any of these solutions (in a particular branch) is non-integer, then that branch of the algorithm terminates since it implies that some solutions of (2.1) have movable algebraic branch point. If any of the resonances are rational, then a change of variables in (2.1) may remove the algebraic branch point. Such changes are not carried out automatically.

For branch (4.11),

$$\begin{aligned} \det(Q_r) &= (r-4)(r-3)^2 \{r^2(1+\beta) - 3r(1+\beta) - 4(1-\beta)\} r^2(r+1) \\ &\quad \times (1+\beta)^5 \bar{u}_0^5(x, t) \bar{v}_0^5(x, t) g_x^8(x, t). \end{aligned} \quad (4.14)$$

Since the roots of (4.14) for r depend on the constant parameter β , we must choose values of β so that all the solutions are integers before proceeding. For $\beta = 1$, the resonances are $r_1 = -1, r_2 = r_3 = r_4 = 0, r_5 = r_6 = r_7 = 3, r_8 = 4$.

While taking $\beta = 0$ also yields all integer resonances, it violates the assumption that all the parameters in (2.1) are nonzero. Allowing the parameters in (2.1) to be zero could cause a false balance in Algorithm 4.1. Thus, (2.1) with $\beta = 0$ should be treated separately. In this example however, setting $\beta = 0$ does not affect the dominant behavior and the resonances are $r_1 = r_2 = -1, r_3 = r_4 = 0, r_5 = r_6 = 3$, and $r_7 = r_8 = 4$.

Although taking $\beta = 25/7$ leads to rational resonances at $r_4 = r_5 = 3/2$, they are not easily resolved by a change of variables in (4.3). The branches with dominant behavior, $\alpha_1 = \alpha_2 = -1, \alpha_3 \geq \alpha_4 \geq 3$, have resonances $r_1 = -\alpha_3 - 1, r_2 = -\alpha_3 + 2, r_3 = -\alpha_4 - 1, r_4 = -\alpha_4 + 2, r_5 = -1, r_6 = 0, r_7 = 3$ and $r_8 = 4$. Since $r_1, r_2 < -1$ when $\alpha_3 = \alpha_4 = 3$, $r_1, r_2, r_3 < -1$ when $\alpha_3 > \alpha_4 = 3$, and $r_1, \dots, r_4 < -1$ when $\alpha_3, \alpha_4 > 3$, these are not principal branches and should be investigated using the perturbative Painlevé approach [9].

4.3 Algorithm to determine the constants of integration and check compatibility conditions

Step 1 (Generate the system for the coefficients of the Laurent series at level k). Substitute

$$u_i(\mathbf{z}) = g^{\alpha_i}(\mathbf{z}) \sum_{k=0}^{r_m} u_{i,k}(\mathbf{z}) g^k(\mathbf{z}) \quad (4.15)$$

into (2.1) and multiply F_i by $g^{-\gamma_i}(\mathbf{z})$, where γ_i is the lowest exponent of $g(\mathbf{z})$ in F_i . The equations for determining the coefficients of the Laurent series at level k then arise by equating to zero the coefficients of $g^k(\mathbf{z})$. These equations, at level k , are linear in $u_{i,k}(\mathbf{z})$ and depend only on $u_{i,j}(\mathbf{z})$ and $g(\mathbf{z})$ (and their derivatives) for $1 \leq i \leq M$ and $0 \leq j < k$. Thus, the system can be written as

$$Q_k \mathbf{u}_k = \mathbf{G}_k(\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{k-1}, g, \mathbf{z}), \quad (4.16)$$

where $\mathbf{u}_k = (u_{1,k}(\mathbf{z}), \dots, u_{M,k}(\mathbf{z}))^T$.

Step 2 (Solve the linear system for the coefficients of the Laurent series). If the rank of Q_k equals the rank of the augmented matrix $(Q_k | \mathbf{G}_k)$, solve (4.16) for the coefficients of the Laurent series. If $k = r_j$, check that $\text{rank } Q_k = M - s_j$, where s_j is the algebraic multiplicity of the resonance r_j in $\det(Q_r) = 0$.

If $\text{rank } Q_k \neq \text{rank}(Q_k | \mathbf{G}_k)$, Gauss reduce the augmented matrix $(Q_k | \mathbf{G}_k)$ to determine the compatibility condition. If all the compatibility conditions can be resolved by restricting the coefficients parameterizing (2.1), then (2.1) has the Painlevé property for those specific values. If any of the compatibility conditions cannot be resolved by restricting the coefficients parameterizing (2.1), then the Laurent series solution for this branch has a movable logarithmic branch point and the algorithm terminates.

For (4.3) with $\beta = 1$, the principal branch

$$\begin{cases} \alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = -1, \\ u_0(x, t) = -\bar{u}_0^{-1}(x, t) \{v_0(x, t) \bar{v}_0(x, t) - 2g_x^2(x, t)\}, \\ \bar{u}_0(x, t), v_0(x, t), \bar{v}_0(x, t) \text{ arbitrary}, \\ r_1 = -1, r_2 = r_3 = r_4 = 0, r_5 = r_6 = r_7 = 3, r_8 = 4, \end{cases} \quad (4.17)$$

has three compatibility conditions at level $k = r_5 = r_6 = r_7 = 3$. These compatibility conditions require that $a_x(x, t) = \bar{a}_x(x, t) = b_x(x, t) = \bar{b}_x(x, t)$ and $c_x(x, t) = d_x(x, t) = 0$.

At level $k = r_8 = 4$, the compatibility condition requires $d(t) = \bar{c}(t)$ and

$$\begin{aligned} & \{(a - \bar{a})^2 + 2i(a_x - \bar{a}_x)h'(t)\}(2 + v_0\bar{v}_0) - \{(b - \bar{b})^2 + 2i(b_x - \bar{b}_x)h'(t)\}v_0\bar{v}_0 \\ & + 2i(a - \bar{a} + b - \bar{b})(v_0(\bar{v}_0)_t + (v_0)_t\bar{v}_0) + i(a_t - \bar{a}_t - b_t + \bar{b}_t)v_0\bar{v}_0 \\ & + 2i(a_t - \bar{a}_t) - (a_{xx} + \bar{a}_{xx} - b_{xx} - \bar{b}_{xx})v_0\bar{v}_0 - 2a_{xx} + 6\bar{a}_{xx} \equiv 0, \end{aligned} \quad (4.18)$$

where we have taken $g(x, t) = x - h(t)$. Careful inspection of (4.18) reveals that $a(x, t) = b(x, t)$. Setting $a(x, t) = b(x, t) = r(x, t) + is(x, t)$, where $r(x, t)$ and $s(x, t)$ are arbitrary real functions, (4.18) becomes

$$2s^2(x, t) + s_t(x, t) + 2h'(t)s_x(x, t) - r_{xx}(x, t) + 2is_{xx}(x, t) \equiv 0. \quad (4.19)$$

Since $h'(t)$ is arbitrary, it follows that $s_x(x, t) = 0$. Thus, $r_{xx}(x, t) = 2s^2(t) + s'(t)$ and upon integration

$$r(x, t) = \frac{1}{2}\{2s^2(t) + s'(t)\}x^2 + r_1(t)x + r_2(t), \quad (4.20)$$

where $r_1(t)$ and $r_2(t)$ are arbitrary functions.

Therefore, the generalized coupled NLS equations,

$$\begin{aligned} iu_t + u_{xx} + (|u|^2 + |v|^2)u + \left\{s^2(t) + \frac{1}{2}s'(t)\right\}x^2 + r_1(t)x + r_2(t) + is(t) u + c(t)v &= 0, \\ iv_t + v_{xx} + (|u|^2 + |v|^2)v + \left\{s^2(t) + \frac{1}{2}s'(t)\right\}x^2 + r_1(t)x + r_2(t) + is(t) v + \bar{c}(t)u &= 0, \end{aligned}$$

passes the Painlevé test, where $r_1(t), r_2(t)$, and $s(t)$ are arbitrary real functions and $c(t)$ is an arbitrary complex function.

When $\beta = 0$, the two compatibility conditions at level $k = r_5 = r_6 = 3$ require that $c(x, t) = d(x, t) = \bar{c}(x, t) = \bar{d}(x, t) = 0$. Similarly, the compatibility conditions at level $k = r_7 = r_8 = 4$, require that

$$a(x, t) = \left\{s^2(t) - \frac{1}{2}s'(t)\right\}x^2 + r_1(t)x + r_2(t) + is(t), \quad (4.21)$$

where $r_1(t), r_2(t)$ and $s(t)$ are arbitrary real functions. Therefore,

$$iu_t + u_{xx} + |u|^2u + \left\{s^2(t) - \frac{1}{2}s'(t)\right\}x^2 + r_1(t)x + r_2(t) + is(t) u = 0, \quad (4.22)$$

passes the Painlevé test, a fact confirmed in [1].

5 Additional Examples

5.1 A peculiar ODE

Consider the ODE [34]

$$u^2u''' - 3(u')^3 = 0. \quad (5.1)$$

Substituting (3.1) into (5.1) gives $\alpha(\alpha + 2)(2\alpha - 1)\chi^3g(z)^{3(\alpha-1)} = 0$. So, both the terms in (5.1) have the same leading exponent, $3(\alpha - 1)$. Using the procedure in Section 4.1, in Step 5 the software automatically runs the test for the default values $\alpha = -3, -2$, and -1 . The choices $\alpha = -1$ and -3 are incompatible with the assumption $u_0 \neq 0$. The leading term vanishes for $\alpha = -2$ and u_0 is arbitrary. Substituting $u(z) = u_0g^{-2}(z) + u_{1,r}g^{r-2}(z)$, we find that $r_1 = -1, r_2 = 0$, and $r_3 = 10$. Thus, the Laurent series solution of (5.1) is

$$u(z) = u_0(z - z_0)^{-2} + u_{10}(z - z_0)^8 + \cdots, \quad (5.2)$$

where z_0, u_0 and u_{10} are arbitrary constants. Hence, (5.1) passes the Painlevé test.

5.2 The sine-Gordon equation

Consider the sine-Gordon equation [1],

$$u_{tt} + u_{xx} = \sin u. \quad (5.3)$$

Using the transformation $v(x, t) = e^{iu(x, t)}$, we obtain a polynomial differential equation

$$vv_{tt} + vv_{xx} - v_t^2 - v_x^2 = \frac{1}{2}v(v^2 - 1). \quad (5.4)$$

The dominant behavior of (5.4) is $v(x, t) \sim 4(g_x^2(x, t) + g_t^2(x, t))g^{-2}(x, t)$, with resonances $r_1 = -1$ and $r_2 = 2$. The Laurent series solution of (5.4) is

$$v = 4(g_x^2 + g_t^2)g^{-2} - 4(g_{xx} + g_{tt})g^{-1} + v_2 + \dots, \quad (5.5)$$

where $g(x, t)$ and $v_2(x, t)$ are arbitrary functions. The sine-Gordon equation passes the Painlevé test and is indeed completely integrable [1].

5.3 The cylindrical Korteweg-de Vries equation

Consider the generalized KdV equation,

$$u_t + 6uu_x + u_{3x} + a(t)u = 0, \quad (5.6)$$

where $a(t)$ is an arbitrary function parameterizing the equation. The dominant behavior of (5.6) is $u(x, t) \sim -2g_x^2(x, t)g^{-2}(x, t)$, with resonances $r_1 = -1, r_2 = 4$ and $r_3 = 6$. At level $k = r_3 = 6$, we obtain the compatibility condition

$$\frac{2a(t)^2 + a'(t)}{6g_x(x, t)} = 0. \quad (5.7)$$

So, (5.6) passes the Painlevé test if $a(t) = \frac{1}{2t}$. In this case, (5.6) reduces to the cylindrical KdV, which is completely integrable as confirmed by other analyses [1].

5.4 A fifth-order generalized Korteweg-de Vries equation

Consider the generalized fifth-order KdV equation,

$$u_t + au_xu_{xx} + buu_{3x} + cu^2u_x + u_{5x} = 0, \quad (5.8)$$

with constant parameters a, b , and c . The dominant behavior of (5.8) is

$$u(x, t) \sim -\frac{3g_x^2(x, t)}{c} \left\{ (a + 2b) \pm \sqrt{a^2 + 4ab + 4b^2 - 40c} \right\} g^{-2}(x, t). \quad (5.9)$$

The resonances are the roots of

$$\begin{aligned} \det(Q_r) = & -c(r - 6)(r + 1) \left(3\sqrt{(a + 2b)^2 - 40c}(2a - b(r - 4)) \right. \\ & \left. - 6(a + 2b)^2 + 240c + (3b(a + 2b) - 86c)r + 15cr^2 - cr^3 \right) g_x^5. \end{aligned} \quad (5.10)$$

Determining what values of a , b , and c that lead to integer roots of (5.10) is difficult by hand or with a computer. An investigation of the scaling properties of (5.8) reveals that only the ratios a/b and c/b^2 are important. Let us consider the well-known special cases.

If we take $a = b$ and $5c = b^2$, then (5.8) passes the Painlevé test with resonances $r_1 = -2, r_2 = -1, r_3 = 5, r_4 = 6, r_5 = 12$ and $r_1 = -1, r_2 = 2, r_3 = 3, r_4 = 6, r_5 = 10$. Taking $b = 5$, equation (5.8) becomes the completely integrable equation

$$u_t + 5u_x u_{xx} + 5uu_{3x} + 5u^2 u_x + u_{5x} = 0, \quad (5.11)$$

due to Sawada and Kotera [36] and Caudrey et al. [5].

If we take $a = 2b$ and $10c = 3b^2$, then (5.8) passes the Painlevé test with resonances $r_1 = -3, r_2 = -1, r_3 = 6, r_4 = 8, r_5 = 10$ and $r_1 = -1, r_2 = 2, r_3 = 5, r_4 = 6, r_5 = 8$. For $b = 10$, equation (5.8) is a member of the completely integrable KdV hierarchy

$$u_t + 10uu_{3x} + 20u_x u_{xx} + 30u^2 u_x + u_{5x} = 0, \quad (5.12)$$

due to Lax [26].

If we take $2a = 5b$ and $5c = b^2$, then (5.8) passes the Painlevé test with resonances $r_1 = -7, r_2 = -1, r_3 = 6, r_4 = 10, r_5 = 12$ and $r_1 = -1, r_2 = 3, r_3 = 5, r_4 = 6, r_5 = 7$. When $b = 10$, equation (5.8) is the Kaup-Kupershmidt equation [12, 17],

$$u_t + 10uu_{3x} + 20u^2 u_x + 25u_x u_{xx} + u_{5x} = 0, \quad (5.13)$$

which is also known to be completely integrable.

While there are many other values for a , b , and c , for which (5.10) only has integer roots, but compatibility conditions prevent (5.8) from having the Painlevé property. For instance, when $a = 2b$ and $5c = 2b^2$, the resonances are $r_1 = -1, r_2 = 0, r_3 = 6, r_4 = 7, r_5 = 8$. At level $k = r_2 = 0$, we are forced to take $u_0(x, t) = -30g_x^2(x, t)/b$, so the Laurent series solution is not the general solution and (5.8) fails the Painlevé test. Similarly, when $7a = 19b$ and $49c = 9b^2$, we have resonances $r_1 = -1, r_2 = 3$ and $r_3 = r_4 = r_5 = 6$, so the Laurent series solution is not the general solution and, again (5.8) fails the Painlevé test.

6 Brief Review of Symbolic Algorithms and Software

There is a variety of methods for testing nonlinear ODEs and PDEs for the Painlevé property. While the WTC algorithm discussed in this paper is the most common method used in Painlevé analysis, it is not appropriate in all cases. For instance, there are numerous completely integrable differential equations which have algebraic branching in their series solutions; a property that is allowed by the so-called “weak” Painlevé test (see [13, 32, 33]). A more thorough approach for testing differential equations with branch points is the poly-Painlevé test (see [22, 23]). The perturbative Painlevé test [9] was developed to check the compatibility conditions of negative resonances other than $r = -1$.

For testing ODEs, there are several implementations: ODEPAINLEVE developed by Rand and Winternitz [34] in *Macysma* is restricted to scalar differential equations; PTEST.RED by Renner in *Reduce* [35]; and, a *Reduce* package by Scheen [37] which implements both the traditional and the perturbative Painlevé tests. For testing PDEs, there are a few implementations. The package PAINMATH.M by Hereman et al. [16] is unable to find all the

dominant behaviors in systems with undetermined α_i and is limited to two independent variables.

Only the *Maple* package `PDEPtest` by Xu and Li [45–47] is comparable to our package `PainleveTest.m` [4]. The package `PDEPtest` was written after our package and allows the testing of systems of PDEs (but not ODEs) parameterized by arbitrary functions using either the traditional WTC algorithm or the simplification proposed by Kruskal (see Section 3). While `PDEPtest` can find all the dominant behaviors in some systems with undetermined α_i (such as the Hirota-Satsuma system), it fails to find the dominant behaviors for systems in which more than one α_i is undetermined (such as the NLS equation, $iu_t + u_{zz} + 2u|u|^2 = 0$, which is completely integrable [1]). Furthermore, `PDEPtest` requires that all the α_i are negative, a weakness of the implementation, since it is standard to allow some positive exponents (see equation (2.4) in [33] with leading exponents -1 and 1).

7 Using the Software Package `PainleveTest.m`

The package `PainleveTest.m` has been tested on both PCs and UNIX work stations with *Mathematica* versions 3.0, 4.0, 4.1, 5.0, 5.1, and 6.0 using a test set of over 50 PDEs and two dozen ODEs. The Backus-Naur form of the function is

```

⟨Main Function⟩ → PainleveTest[⟨Equations⟩,⟨Functions⟩,
                               ⟨Variables⟩,⟨Options⟩]
⟨Options⟩ → Verbose → ⟨Boolean⟩ |
            KruskalSimplification → ⟨Variable⟩ |
            DominantBehaviorMin → ⟨Negative Integer⟩ |
            DominantBehaviorMax → ⟨Integer⟩ |
            DominantBehavior → ⟨List of Rules⟩ |
            DominantBehaviorConstraints → ⟨List of Constraints⟩ |
            DominantBehaviorVerbose → ⟨Range⟩ |
            ResonancesVerbose → ⟨Range⟩ |
            ConstantsOfIntegrationVerbose → ⟨Range⟩
⟨Bool⟩ → True | False
⟨Range⟩ → 0 | 1 | 2 | 3
⟨List of Rules⟩ → {{alpha[1] → ⟨Integer⟩, alpha[2] → ⟨Integer⟩, ...}, ...}
⟨List of Constraints⟩ → {alpha[1] == alpha[2], ...}

```

The output of the function is

```

{ { {Dominant behavior}, {Resonances},
  { {Laurent series coefficients}, {Compatibility conditions}, ... } } }

```

If using a PC, place the package `PainleveTest.m` in a directory, say `myDirectory` on drive C. Start a *Mathematica* notebook session and execute the commands:

```
In[1] = SetDirectory["c:\\myDirectory"]; (* Specify the directory *)
```

```
In[2] = Get["PainleveTest.m"] (* Read in the package *)

In[3] = PainleveTest[ (* Test the KdV equation *)
  {D[u[x,t],t]+6*u[x,t]*D[u[x,t],x]+D[u[x,t],{x,3}] == 0},
  u[x, t], {x,t}, KruskalSimplification -> x]
```

```
Out[3] =
```

$$\left\{ \left\{ \left\{ \alpha_1 \rightarrow -2 \right\}, \left\{ r \rightarrow -1, r \rightarrow 4, r \rightarrow 6 \right\}, \right. \right. \\ \left. \left\{ \left\{ u_{1,0} \rightarrow -2, u_{1,1} \rightarrow 0, u_{1,2} \rightarrow \frac{h'(t)}{6}, u_{1,3} \rightarrow 0, \right. \right. \right. \\ \left. \left. \left. u_{1,4} \rightarrow C_1(t), u_{1,5} \rightarrow \frac{h''(t)}{36}, u_{1,6} \rightarrow C_2(t) \right\}, \left\{ \right\} \right\} \right\}$$

The option `KruskalSimplification -> x` allows one to use $g(x,t) = x - h(t)$ in the calculation of the constants of integration and in checking the compatibility conditions.

```
In[4] = PainleveTest[ (* Eq. (2.4) in Ramani et al. [32] *)
  {D[x[z], z] == x[z]*(a - x[z] - y[z]),
  D[y[z], z] == y[z]*(x[z] - 1)},
  {x[z], y[z]}, {z}, DominantBehaviorMax -> 1 ]
```

```
Out[4] =
```

$$\left\{ \left\{ \left\{ \alpha_1 \rightarrow -1, \alpha_2 \rightarrow -1 \right\}, \left\{ r \rightarrow -1, r \rightarrow 2 \right\}, \left\{ \left\{ u_{1,0} \rightarrow -1, \dots \right\}, \left\{ a + 1 = 0 \right\} \right\} \right\}, \right. \\ \left. \left\{ \left\{ \alpha_1 \rightarrow -1, \alpha_2 \rightarrow 1 \right\}, \left\{ r \rightarrow -1, \rightarrow 0 \right\}, \left\{ \left\{ u_{1,0} \rightarrow 1, u_{2,0} \rightarrow C_1 \right\}, \left\{ \right\} \right\} \right\} \right\}$$

In this example, if the `DominantBehaviorMax` option was not used, we would wrongly conclude that the system only passes the Painlevé test when $a = -1$. However, by allowing positive α_i , we find the second branch $\alpha_1 = -1$ and $\alpha_2 = 1$, for which the system passes the Painlevé test without restricting the value of the parameter a . Alternatively, executing

```
In[5] = PainleveTest[{ D[x[z], z] == x[z]*(a - x[z] - y[z]),
  D[y[z], z] == y[z]*(x[z] - 1)}, {x[z], y[z]}, {z},
  DominantBehavior -> {{alpha[1] -> -1, alpha[2] -> 1}} ]
```

would only test the branch with $\alpha_1 = -1$ and $\alpha_2 = 1$. For an example of the option `DominantBehaviorConstraints`, see Step 4 of Algorithm 4.1.

The option `Verbose -> True` gives a brief trace of the calculations in each of the three steps of the algorithm. The options `DominantBehaviorVerbose`, `ResonancesVerbose`, and `ConstantsOfIntegrationVerbose` allow for a more detailed trace of the calculation. For instance, `DominantBehaviorVerbose -> 1` would show the result of substituting the ansatz, the exponents before and after removing non-dominant powers, etc. While `DominantBehaviorVerbose -> 3` shows the result of nearly every line of code in the package, allowing the user to check the results in the trickiest cases.

8 Discussion and Conclusions

Our software package `PainleveTest.m` is applicable to polynomial systems of nonlinear ODEs and PDEs. While the Painlevé test does not guarantee complete integrability, it helps in identifying candidate differential equations for complete integrability in a straightforward manner. For differential equations with parameters (including arbitrary functions of the independent variables), our software allows the user to determine the conditions under which the differential equations may possess the Painlevé property. Therefore, by finding the compatibility conditions, classes of parameterized differential equations can be analyzed and candidates for complete integrability can be identified.

The difficulty in completely automating the Painlevé test lies in determining the dominant behaviors of the Laurent series solutions; specifically, determining *all* the valid dominant behaviors when one or more of the α_i are undetermined. While there are other implementations for the Painlevé test, ours is currently the only implementation in *Mathematica* which allows the testing of polynomial systems of nonlinear PDEs with no limitations on the number of differential equations or the number of independent variables (except where limited by memory).

References

- [1] Ablowitz M J and Clarkson P A, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, U.K., 1991.
- [2] Ablowitz M J, Ramani A, and Segur H, A Connection Between Nonlinear Evolution Equations and Ordinary Differential Equations of P -type. I, *J. Math. Phys.* **21** (1980), 715–721.
- [3] Ablowitz M J, Ramani A, and Segur H, A Connection Between Nonlinear Evolution Equations and Ordinary Differential Equations of P -type. II, *J. Math. Phys.* **21** (1980), 1006–1015.
- [4] Baldwin D and Hereman W, `PainleveTest.m`: A *Mathematica* Package for the Painlevé Test of Systems of Nonlinear Ordinary and Partial Differential Equations, 2001; Available at http://www.mines.edu/fs_home/whereman/software/painleve/mathematica.
- [5] Caudrey P J, Dodd R K, and Gibbon J D, A New Hierarchy of Korteweg-de Vries Equations, *Proc. Roy. Soc. Lond. A* **351** (1976), 407–422.
- [6] Clarkson P A, The Painlevé Property and a Partial Differential Equations with an Essential Singularity, *Phys. Lett. A* **109** (1985), 205–208.
- [7] Conte R, Singularities of Differential Equations and Integrability, in *Introduction to Methods of Complex Analysis and Geometry for Classical Mechanics and Non-Linear Waves*, Editors: Benest D and Froeschlé C, Gif-sur-Yvette: Editions Frontières, 1993, 49–143.
- [8] Conte R, editor, *The Painlevé Property: One Century Later*, CRM Series in Mathematical Physics, Springer Verlag, New York, 1999.
- [9] Conte R, Fordy A P, and Pickering A, A Perturbative Painlevé Approach to Nonlinear Differential Equations, *Physica D* **69** (1993), 33–58.
- [10] Ercolani N and Siggia E D, Painlevé Property and Integrability, in *What is Integrability?*, Editor: Zakharov V E, *Springer Series in Nonlinear Dynamics*, Springer Verlag, New York, 1991, 63–72.
- [11] Fokas A S, Symmetries and Integrability, *Stud. Appl. Math.* **77** (1987), 253–299.

-
- [12] Fordy A P and Gibbons J, Some Remarkable Nonlinear Transformations, *Phys. Lett. A* **75** (1980), 325–325.
- [13] Goriely A, Integrability and Nonintegrability of Dynamical Systems, *Advanced Series in Nonlinear Dynamics* **19**, World Scientific Publishing Company, Singapore, 2001.
- [14] Grammaticos B and Ramani A, Integrability – and How to Detect It, in Integrability of Nonlinear Systems, Editors: Kosmann-Schwarzbach Y, Grammaticos B, and Tamizhmani K, Springer Verlag, Berlin, 1997, 30–94.
- [15] Hereman W, Banerjee P P, and Chatterjee M, Derivation and Implicit Solutions of the Harry Dym Equation, and Its Connections with the Korteweg-de Vries Equation, *J. Phys. A* **22** (1989), 241–255.
- [16] Hereman W, Göktaş Ü, Colagrosso M D, and Miller A J, Algorithmic Integrability Tests for Nonlinear Differential and Lattice Equations, *Comp. Phys. Comm.* **115** (1998), 428–446.
- [17] Hirota R and Ramani A, The Miura Transformations of Kaup’s Equation and of Mikhailov’s Equation, *Phys. Lett. A* **76** (1980), 95–96.
- [18] Hone A, Painlevé Tests, Singularity Structure, and Integrability (2005), 1–34, Preprint UKC/IMS/03/33, Institute of Mathematics, Statistics, and Actuarial Science, University of Kent, Canterbury, U.K.; [arXiv:nlin.SI/0502017](https://arxiv.org/abs/nlin.SI/0502017).
- [19] Ince E L, Ordinary Differential Equations, Dover Publishing Company, New York, 1944.
- [20] Jimbo M, Kruskal M D, and Miwa T, Painlevé Test for the Self-Dual Yang-Mills Equation, *Phys. Lett. A* **92** (1982), 59–60.
- [21] Johnson R S, On Solutions of the Camassa-Holm Equation, *Proc. Roy. Soc. Lond. A* **459** (2003), 1687–1708.
- [22] Kruskal M D and Clarkson P A, The Painlevé-Kowalevski and poly-Painlevé Tests for Integrability, *Stud. Appl. Math.* **86** (1992), 87–165.
- [23] Kruskal M D, Joshi N, and Halburd R, Analytic and Asymptotic Methods for Nonlinear Singularity Analysis: A Review and Extensions of Tests for the Painlevé Property, in Proceedings of CIMPA Summer School on Nonlinear Systems, Editors: Grammaticos B and Tamizhmani K, *Lecture Notes in Physics* **495**, Springer Verlag, Heidelberg, 1997, 171–205.
- [24] Lakshmanan M and Kaliappan P, Lie Transformations, Nonlinear Evolution Equations, and Painlevé Forms, *J. Math. Phys.* **24** (1983), 795–806.
- [25] Lamb G L, Elements of Soliton Theory, Wiley and Sons, New York, 1980.
- [26] Lax P D, Integrals of Nonlinear Equations of Evolution and Solitary Waves, *Comm. Pure Appl. Math.* **21** (1968), 467–490.
- [27] McLeod J B and Olver P J, The Connection Between Partial Differential Equations Soluble by Inverse Scattering and Ordinary Differential Equations of Painlevé Type, *SIAM J. Math. Anal.* **14** (1983), 488–506.
- [28] Newell A C, Tabor M, and Zeng Y B, A Unified Approach to Painlevé Expansions, *Physica D* **29** (1987), 1–68.
- [29] Osgood W F, Topics in the Theory of Functions of Several Complex Variables, in The Madison Colloquium 1913, *Colloquium Lectures* **4**, AMS, New York, 1914, 111–230.
- [30] Painlevé P, Mémoire sur les Equations Différentielles dont l’Intégrale Générale est Uniforme, *Bull. Soc. Math. France* **28** (1900), 201–261.
- [31] Pickering A, The Singular Manifold Method Revisited, *J. Math. Phys.* **37** (1996), 1894–1927.

-
- [32] Ramani A, Dorizzi B, and Grammaticos B, Painlevé Conjecture Revisited, *Phys. Rev. Lett.* **49** (1982), 1539–1541.
- [33] Ramani A, Grammaticos B, and Bountis T, The Painlevé Property and Singularity Analysis of Integrable and Nonintegrable Systems, *Phys. Rep.* **180** (1989), 159–245.
- [34] Rand D W and Winternitz P, ODEPAINLEVE – A MACSYMA Package for Painlevé Analysis of Ordinary Differential Equations, *Comp. Phys. Comm.* **42** (1986), 359–383.
- [35] Renner F, A Constructive REDUCE Package Based Upon the Painlevé Analysis of Nonlinear Evolutions Equations in Hamiltonian and/or Normal Form, *Comp. Phys. Comm.* **70** (1992), 409–416.
- [36] Sawada S and Kotera T, A Method for Finding N-Soliton Solutions of the KdV and KdV-Like Equation, *Prog. Theor. Phys.* **51** (1974), 1355–1367.
- [37] Scheen C, Implementation of the Painlevé Test for Ordinary Differential Equations, *Theor. Comp. Sci.* **187** (1997), 87–104.
- [38] Steeb W H and Euler N, Nonlinear Evolution Equations and Painlevé Test, World Scientific Publishing Company, Singapore, 1988.
- [39] Tabor M, Painlevé Property for Partial Differential Equations, in Soliton Theory: A Survey of Results, Editor: Fordy A P, Manchester University Press, Manchester, U.K., 1990, 427–446.
- [40] Tan Y and Yang J, Complexity and Regularity of Vector-Soliton Collisions, *Phys. Rev. E* **64** (2001), 056616.
- [41] Ward R S, The Painlevé Property for the Self-dual Gauge-field Equations, *Phys. Lett. A* **102** (1984), 279–282.
- [42] Weiss J, The Painlevé Property for Partial Differential Equations. II: Bäcklund Transformations, Lax Pairs, and the Schwarzian Derivative, *J. Math. Phys.* **24** (1983), 1405–1413.
- [43] Weiss J, Bäcklund Transformation and Linearizations of the Hénon-Heiles System, *Phys. Lett. A* **102** (1984), 329–331.
- [44] Weiss J, Tabor M, and Carnevale G, The Painlevé Property for Partial Differential Equations, *J. Math. Phys.* **24** (1983), 522–526.
- [45] Xu G Q and Li Z B, A Maple Package for the Painlevé Test of Nonlinear Partial Differential Equations, *Chin. Phys. Lett.* **20** (2003), 975–978.
- [46] Xu G Q and Li Z B, Symbolic Computation of the Painlevé Test for Nonlinear Partial Differential Equations Using Maple, *Comp. Phys. Comm.* **161** (2004), 65–75.
- [47] Xu G Q and Li Z B, PDEPtest: A Package for the Painlevé Test of Nonlinear Partial Differential Equations, *Appl. Math. Comp.* **169** (2005), 1364–1379.
- [48] Yoshida H, Necessary Condition for the Existence of Algebraic First Integrals I & II, *Celes. Mech.* **31** (1983), 363–399.
- [49] Ziglin S L, Self-Intersection of the Complex Separatrices and the Nonexistence of the Integrals in the Hamiltonian Systems with One-and-half Degrees of Freedom, *J. Appl. Math. Mech.* **45** (1982), 411–413.
- [50] Ziglin S L, Branching of Solutions and Nonexistence of First Integrals in Hamiltonian Mechanics I, *Func. Anal. Appl.* **16** (1983), 181–189.
- [51] Ziglin S L, Branching of Solutions and Nonexistence of First Integrals in Hamiltonian Mechanics II, *Func. Anal. Appl.* **17** (1983), 6–17.